

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : A61K 7/08, C11D 1/12, 1/94	A1	(11) International Publication Number: WO 92/08440 (43) International Publication Date: 29 May 1992 (29.05.92)
(21) International Application Number: PCT/GB91/01955 (22) International Filing Date: 7 November 1991 (07.11.91) (30) Priority data: 9024162.1 7 November 1990 (07.11.90) GB (71) Applicant (for AU CA GB only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (71) Applicant (for all designated States except AU CA GB): UNILEVER NV [NL/NL]; Burgemeester 's Jacobplein 1, NL-Rotterdam (NL). (72) Inventors: LEE, Robert, Stanley ; 37 Poulton Road, Spital, Wirral L63 9LD (GB). SALMON, Tom, Matthew, Forrest ; 48 Walpole Street, Chester CH1 4HG (GB).		(74) Agent: FORD, M., F.; Mewburn Ellis, 2 Cursitor Street, London EC4A 1BQ (GB). (81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: DETERGENT COMPOSITION (57) Abstract <p>Aqueous detergent compositions giving good mildness, good lather volume and lather creaminess contain acyl isethionate, a zwitterionic detergent which is a quaternised amino acid, usually a betaine, and another anionic detergent. Proportions and ratios are specified and alkanolamides are absent or limited in amount. Incorporation of the additional anionic detergent, such as ether sulphate, has surprisingly little adverse effect and provides economy.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU ⁺	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE ⁺	Germany	MC	Monaco	US	United States of America
DK	Denmark				

⁺ Any designation of "SU" has effect in the Russian Federation. It is not yet known whether any such designation has effect in other States of the former Soviet Union.

DETERGENT COMPOSITION

This invention relates to detergent compositions in liquid or gel form, suitable for personal washing of either skin or hair. It is desirable that such compositions should be mild to the skin (even if intended for use as a hair shampoo) yet able to generate lather which the user will judge to be good, both in quantity and quality. It is not easy to achieve mildness simultaneously with good lather. For example sodium lauryl sulphate is high foaming but harsh while alcohol ethoxylates are mild but low foaming. It is of course also desirable to achieve economy in cost of materials.


We have now found that a good combination of properties can be obtained by use of a detergent mixture in which a combination of fatty acyl isethionate and a zwitterionic detergent constitute a high proportion of the detergent mixture, but some other anionic detergent is also included.

Some prior proposals for compositions containing fatty acyl isethionate have included polar nonionic detergents, notably alkanolamides such as coconut mono- or di-ethanolamide. This is logical because these materials are well known as lather enhancers. Surprisingly we have found that alkanolamides are not helpful in combinations of acyl isethionate and betaine, and their use has undesirable

effects.

According to the present invention there is provided a detergent composition in the form of an aqueous liquid or gel, comprising 10 to 50% by weight of a detergent mixture which comprises

(a) 10 to 60% by weight of the detergent mixture of a fatty acyl isethionate of formula



10 where R is an alkyl group of 7 to 21 carbon atoms and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium,

(b) 10 to 80% by weight of the detergent mixture of a zwitterionic detergent which has a hydrophilic head group
15 containing a quaternary nitrogen atom and at least one acid group,

(c) 10 to 55% by weight of the detergent mixture of a further anionic detergent,

wherein the amount by weight of the fatty acyl
20 isethionate (a) is not more than three times the amount by weight of the zwitterionic detergent (b), the total of (a) and (b) is from 45 to 90% by weight of the detergent composition, and the composition is sufficiently free of alkanolamide detergents that the amount by weight of
25 alkanolamide is not more than one quarter the amount of the zwitterionic detergent (b).

The materials used in the present application, and the

relationships between them will now be discussed in more detail.

Fatty acyl isethionates may be prepared by the reaction between alkali metal isethionate and aliphatic fatty acids (or their acid chlorides) having from 8 to 22 carbon atoms. Preferably these fatty acids have an iodine value of less than 20. Generally a mixture of aliphatic fatty acids will be used. In one embodiment of the invention at least three quarters of the fatty acyl groups in the acyl isethionate have from 12 to 18 carbon atoms while the balance, up to a quarter of the fatty acyl groups, may have from 8 to 10 carbon atoms. Notably the fatty acyl groups may be provided by coconut fatty acid.

15

We have found that fatty acyl isethionate contributes to mildness in the detergent mixtures of this invention, and also contributes to obtaining foam of good volume and/or quality, as will be mentioned again below. Even fairly low levels of acyl isethionate will contribute to these benefits.

A difficulty with fatty acyl isethionate is that it has a low solubility in water which is an obstacle to incorporating it into aqueous compositions. Typical solubility in distilled water is 0.01% by weight at 25°C. We have found that the use of the zwitterionic detergent required for this invention can achieve dissolution of acyl

isethionate at a much greater concentration than would dissolve in the absence of this second detergent.

The difficulty of solubilising isethionate is of increasing significance as the percentage of this material in the composition increases, of course. In certain preferred forms of the present invention the concentration of fatty acyl isethionate is at least 6% by weight of the composition. Dissolution of the acyl isethionate in an aqueous phase is desirable, since it can lead to a product which is more attractive to the consumer and more stable during storage. It is also valuable in that it simplifies the manufacturing process. Consequently in preferred forms of the compositions the detergent mixture is fully soluble in the aqueous phase of the composition. The composition may then be an isotropic solution or may lack optical clarity solely because of some other constituent such as an opacifying or pearlescent agent.

Materials which do not readily enter aqueous solution include free fatty acids, especially those of longer chain length. It is preferred that fatty acids, especially those with 16 or more carbon atoms, are absent from compositions of this invention.

For solubilising fatty acyl isethionate it is generally desirable that the quantity of the zwitterionic detergent is not less than one third the weight of fatty

acyl isethionate. Better is that the amount of zwitterionic detergent is at least half the weight of isethionate.


5 Use of fatty acyl isethionate jointly with zwitterionic detergent is also advantageous in yielding good foam volume and/or foam with a thick, creamy quality. Both of these properties are important to the end user's perception of the quality of the product. One or other of
10 these advantageous properties can be obtained even when the proportion of zwitterionic is relatively high. However, the quality and/or quantity of foam is generally best when the ratio of fatty acyl isethionate to zwitterionic
15 detergent lies in a range from 3:1 to 1:3. A range from 2:1 to 1:2 is preferred.

 The zwitterionic detergent, when used jointly with isethionate, enhances mildness. For particularly good mildness it is preferred that the zwitterionic detergent is
20 at least 30% by weight of the mixture of detergents. For good lather, it is preferred not to have much more than 60% by weight of zwitterionic. This is consistent with the above preferred range of isethionate to zwitterionic.

25 The combined amount of zwitterionic and acyl isethionate detergents may well be at least 50 or 55% of the detergent mixture, for instance 55 to 75% for the sake of mildness together with economy from inclusion of other

anionic detergent. The range from 75 to 90% of the detergent mixture can also be utilised, however.

Zwitterionic detergents for use in this invention include at least one acid group. This may be a carboxylic or a sulphonic acid group. They include quaternary nitrogen and therefore are quaternary amino acids. They should generally include an alkyl or alkenyl group of 7 to 18 carbon atoms. They will usually comply with an overall structural formula



where R^1 is alkyl or alkenyl of 7 to 18 carbon atoms
 R^2 and R^3 are each independently alkyl, hydroxyalkyl or carboxyalkyl of 1 to 3 carbon atoms


m is 2 to 4

n is 0 or 1


X is alkylene of 1 to 3 carbon atoms optionally substituted with hydroxyl, and

Y is $-CO_2^-$ or $-SO_3^-$

Suitable zwitterionic detergents within the above general formula include simple betaines of formula:



and amido betaines of formula:




where m is 2 or 3.


10

In both formulae R^1 , R^2 and R^3 are as defined previously. R^1 may in particular be a mixture of C_{12} and C_{14} alkyl groups derived from coconut so that at least half, preferably at least three quarters of the groups R^1 have 10 to 14 carbon atoms. R^2 and R^3 are preferably methyl.

A further possibility is that the zwitterionic detergent is a sulphobetaine of formula




25 or



30

where m is 2 or 3, or variants of these in which $-(\text{CH}_2)_3 \text{SO}_3^-$ is replaced by



35

In these formulae R^1 , R^2 and R^3 are as discussed previously.

The invention also requires that some anionic
5 detergent other than acyl isethionate is included in the composition. Conventional anionic detergents are not effective to solubilise fatty acyl isethionate, so it could not be predicted that their presence would be useful or even tolerable. However, we have found that some anionic
10 detergent can be included as a proportion of the detergent mixture, with surprisingly little detriment to the mildness of the composition. Incorporation of anionic detergent is beneficial in that it can act as a partial replacement for the mixture of acyl isethionate and
15 zwitterionic. This can give a saving in cost because acyl isethionate and zwitterionic detergent are both relatively expensive materials.

A further advantage for the addition of a second
20 anionic detergent is in providing greater control over viscosity. Binary mixtures of acyl isethionate and zwitterionic detergents often have higher low-shear viscosity than desired for some product formulations. While viscosity can be reduced by the addition of alcohol
25 or polyol hydrotropes, lather performance is reduced by the inclusion of these materials. The use of other anionic detergent allows the systems to be formulated with control of viscosity in the desired range by addition of

electrolyte and with surprisingly little reduction in lathering characteristics.

The anionic detergent which is included is particularly envisaged as ether sulphate of the formula

$$R^4 O(CH_2 CH_2 O)_t SO_3 M$$

where R^4 is alkyl or alkenyl of 8 to 18 carbon atoms, especially 11 to 15 carbon atoms, t has an average value of at least 2.0 and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium. Preferably t has an average value of 3 or more.

Other anionic detergents may be used. Possibilities include alkyl glyceryl ether sulphates, sulphosuccinates, taurates, sarcosinates, sulphoacetates, alkyl phosphates and acyl lactates. Sulphosuccinates may be monoalkyl sulphosuccinates having the formula:

$R^5 O_2 CCH_2 CH(SO_3 M)CO_2 M$; and amido-MEA sulphosuccinates of the formula: $R^5 CONHCH_2 CH_2 O_2 CCH_2 CH(SO_3 M)CO_2 M$; wherein R^5 ranges from $C_8 - C_{20}$ alkyl, preferably $C_{12} - C_{15}$ alkyl and M is a solubilising cation.

Sarcosinates are generally indicated by the formula $R^5 CON(CH_3)CH_2 CO_2 M$, wherein R ranges from $C_8 - C_{20}$ alkyl, preferably $C_{12} - C_{15}$ alkyl and M is a solubilising cation.

Taurates are generally identified by the formula $R^5 CONR^6 CH_2 CH_2 SO_3 M$, wherein R^5 ranges from $C_8 - C_{20}$ alkyl,

preferably $C_{12}-C_{15}$ alkyl, R^6 ranges from C_1-C_4 alkyl, and M is a solubilising cation.

The anionic detergent included in the composition will generally be selected to avoid harsh detergent such as primary alkane sulphonate or alkyl benzene sulphonate. The amount, if any, of these is preferably less than 3% of the detergents present.

The anionic detergent or mixture of anionic detergents is preferably sufficiently mild in its own right that if tested alone by the zein solubilisation test described in Example 2 below, it causes no greater solubilisation than does an equal concentration of sodium lauryl ether sulphate with average two ethylene oxides per molecule.

Alkanolamide detergents are required to be included at only a low level, if at all. We have found that they reduce mildness rather considerably, even if used in a mixture with the specified zwitterionic detergent. Preferably they are restricted to not more than 5% by weight of the detergent mixture, or one quarter the amount of the zwitterionic, whichever is less. Even better is to exclude alkanolamides and the harsh anionics, alkyl benzene sulphonate and primary alkane sulphonate completely.

Provided the requirements for the minimum quantities of fatty acyl isethionate, the zwitterionic detergent and

other anionic detergent are met, additional amphoteric or nonionic detergent may be included. Preferred, however, is to avoid amine oxide, which reduces lather quality. Accordingly, if amine oxide is present at all, the amount
5 may also be less than 5% of the detergents present.

Other materials may be included in compositions of this invention. Possibilities include colouring agents, opacifying agents, organic polymers, perfumes including
10 deodorant perfumes, bactericidal agents to reduce the microflora on skin, antioxidants and other preservatives, and skin feel modifiers.

Organic polymers which may be present include cross-
15 linked polyacrylates such as the Carbopol polymers available from Goodrich. These can function to increase viscosity or enhance stability of a composition. Polysaccharides are also well known as thickening agents and many are cellulose or cellulose derivatives.

20

The compositions of this invention will generally be pourable liquids or semi-liquids, although they may be somewhat viscous. For this, they may be thickened by including electrolyte. Ammonium salts are preferred over
25 sodium salts which reduce the solubility of fatty acyl isethionate.

Compositions of this invention may be formulated as

products for washing the skin, e.g. bath or shower gels, hand washing compositions or facial washing liquids. They may also be formulated as hair shampoos.


5 A viscosity of at least 0.5 Pa.sec at a low shear rate of 10 to 25 sec⁻¹ may be desired for most product forms. Hair shampoos will generally have a viscosity of at least 1 Pa.sec at such shear rate. Products for washing the skin are customarily more viscous. A viscosity of at least 2 or
10 even at least 3 Pa.sec at the same shear rate is usually appropriate for these.

Example 1

Various binary mixtures were made containing sodium
15 cocoyl isethionate and another detergent. The sodium cocoyl isethionate was Fenopon (Registered Trade Mark) AC78 available from GAF Corporation. It has a water solubility, in distilled water at 25°C, of about 0.01% by weight.


20 Other detergents were:

(1) coconut amidopropyl betaine (CAPB) of formula:




in which R¹CO- is a mixture of acyl groups derived
30 from coconut with C₁₂ and C₁₄ together constituting a majority. This was Tegobetain L7 from Goldschmidt.

(ii) coconut betaine of formula:



in which $\text{R}^1 \text{CO}-$ is a mixture of acyl groups derived from coconut. This betaine was Empigen BB from Albright and Wilson.

(iii) coconut amidopropyl sulphobetaine of formula:



in which $\text{R}^1 \text{CO}$ is again coconut-derived acyl. This was Rewoteric AM-CAS from Rewo.

(iv) coconut diethanolamide (CDEA) which was Empilan CDE from Albright and Wilson.

(v) sodium lauryl ether sulphate with average three ethylene oxide residues (SLES) which was Empicol 0251 from Albright and Wilson.

(vi) disodium lauryl ether sulphosuccinate with average 3EO, which was Rewopol SBFA from Rewo.

25

The amount of mixture which could be dissolved to an isotropic solution in demineralized water was measured for mixtures containing 80%, 60% and 40% of the isethionate, with 20%, 40% and 60% respectively of each other
30 detergent.

The results are set out in the following Table 1.

Table 1

Solubilities of binary mixtures as isotropic
solutions in distilled water (% w/w)

5	Second detergent in mixture	Ratio of isethionate : second detergent		
		80:20	60:40	40:60
10	(i) CAPB	below 10%	15-20%	35-40%
	(ii) betaine	15-20%	30-35%	30-35%
	(iii) sulpho- betaine	below 10%	20-25%	25-30%
15	(iv) CDEA	below 10%	30-40%	30-40%
	(v) SLES	below 10%	below 10%	below 10%
20	(vi) sulpho- succinate	below 10%	below 10%	below 10%

This shows that the zwitterionic detergent (i), (ii)
and (iii) were effective to solublise the acyl isethionate,
25 unlike the two anionic detergents (v) and (vi). CDEA was
also effective.

Example 2

A number of detergents and mixtures were assessed for
30 mildness using the zein solubilisation test.

This test was first described by Gotte, Proc. Int.
Congr. Surface Active Subs., 4th, Brussels 3 83-90, 1094.
In this test, as carried out by us, 5g of zein (which is a
35 protein available from Kodak) was mixed with 40ml of a
detergent solution (which in this example contained 1.2g
detergent), then shaken for 1 hour at 35°C. Solids were

then removed by centrifuging. The supernatant was filtered and the amount of protein in the filtered supernatant was determined by analysis for nitrogen. Correction was made for any nitrogen from the detergent itself. The result is a measure of the harshness of the detergent, since a mild detergent solubilises protein from the zein to a lesser extent than a harsh detergent.

The detergents were sodium cocoyl isethionate (Fenopon AC78 as in Example 1), cocoamidopropyl betaine (CAPB) which was Rewoteric AMB14 from Rewo and sodium lauryl ether sulphate (SLES) with average 3EO (Genapol ZRO from Hoechst). All materials and mixtures were tested at an overall detergent concentration of 3% by weight.

15

All the detergent solutions were clear, isotropic liquids. The proportions of detergents in the solutions tested, and the zein solubilisation results are set out in the following Table:

20

Table 2

		<u>wt% of total detergent</u>									
Example no.		2A	2B	2C	2D	2E	2F	2G	2H	2J	2K
	Isethionate:	0	0	18	0	50	50	16	33	15	30
25	CAPB:	0	33	18	20	25	40	42	33	70	15
	SLES:	100	66	64	80	25	10	42	33	15	55
zein solubilisation:											
(%N)		0.55	0.35	0.25	0.40	0.24	0.15	0.07	0.19	0.09	0.25

Comparison of Examples 2A, 2B and 2D shows that the CAPB improves mildness of SLES. However, Example 2C is milder than Example 2B with the same amount of SLES and
5 also milder than Example 2D with the same amount of CAPB.

Examples 2E to 2K show good mildness over the area where isethionate and CAPB together constitute at least 45% by weight of the detergent active, especially when the CAPB
10 is at least 30% by weight of the total detergent.

Example 3

The foaming properties of various compositions were assessed by a panel of twenty persons trained in magnitude
15 estimation techniques. Each panellist wore surgical gloves which had first been washed with soap to remove any talc.

0.5g of test composition was dosed onto the wet gloves, and the panellist rubbed his or her hands together
20 to generate lather. The panellist estimated the magnitude of the volume of lather and also the extent to which the lather appears thick and creamy.

The results are expressed as the normalised averages
25 of the scores given by the panellists. We have found good correlation between estimates of lather volume and volumes actually collected into a measuring cylinder.

In this example the compositions were isotropic aqueous solutions of sodium cocoyl isethionate (Fenopon AC78 as in Example 1) and a simple betaine. (Empigen BB as in Example 1).

5

A control solution contained sodium lauryl ether sulphate (average 3EO) and the same betaine (a mixture which is typical of a conventional shower gel or shampoo).

10 All the solutions contained 15% by weight of detergent. The results obtained were:

Table 3

panel scores

15	Weight ratio		lather volume	creaminess
	isethionate	: betaine		
	70	: 30	139	145
	60	: 40	110	120
	50	: 50	110	120
20	40	: 60	88	121
	Control			
	SLES	: betaine		
	86	: 14	101	87

25 This shows that the isethionate mixtures gave better foam volume than the control solution except when the quantity of betaine exceeded the quantity of isethionate, while the creaminess of the lather was better at all the proportions of betaine.

Example 4

The procedure of Example 3 was repeated using solutions of cocoyl isethionate, betaine and sodium lauryl ether sulphate, with average 3EO. These were Fenopon AC78, Empigen BB and Empicol 0251 all as in Example 1. The same control solution of SLES and betaine was used. All solutions were isotropic and contained a total of 15% by weight of detergent. The results obtained were:

10

Table 4

Weight ratio				<u>panel scores</u>	
isethionate : betaine : SLES				lather volume	lather
15	creaminess				
	60	30	10	112	99
	60	20	20	96	99
	50	40	10	105	100
	50	20	30	97	100
20	40	30	30	97	102
	30	35	35	94	99
		14	86	98	97

It will be observed that lather volume and creaminess were about equal to the control or better.

25

Example 5

A number of aqueous solutions of sodium cocoyl isethionate and a second detergent were made and assessed

for lathering as in Example 3. The isethionate was Fenopon AC78 as in Examples 1 and 3. The second detergent was either Empigen BB as in Examples 1 and 3, or a C_{12} to C_{14} alkyl dimethyl amine oxide (Empigen OB from Albright and Wilson). The same ether sulphate/betaine control solution was used. All solutions contained 15% by weight of detergent. The results obtained were:

Table 5

10	Weight ratio			<u>panel scores</u>	
	isethionate : creaminess	betaine :	amine oxide	lather volume	
	50	--	50	115	88
15	60	--	40	115	95
	60	40	--	100	110
	70	--	30	97	100
	70	30	--	105	90
	Control	86 SLES:	14 betaine	85	99

20

This shows that amine oxide gave some superiority in foam volume, when used at higher proportions but was then inferior to betaine in creaminess of lather.

25 Example 6

Solutions were prepared containing sodium cocoyl isethionate (Fenopon AC78 as in Example 1) sodium lauryl ether sulphate with average 3EO and another detergent active which was either or both of coconut diethanolamide

(CDEA) and coconut amidopropyl betaine (CAPB: Rewoteric AMB14 as in Example 2).

5 All solutions contained 15% by weight detergent. The amounts of detergent, as weight percentages of total detergent, together with the total panel scores for lather and creaminess were:

Table 6

10	<u>% by weight based on total detergent</u>					
	Isethionate	33	33	33	50	50
	CAPB	54	-	27	40	-
	CDEA	-	54	27	-	40
	SLES	13	13	13	10	10
15		<u>panel scores</u>				
	Volume	125	86	98	105	92
	Creaminess	114	99	110	92	86

20 These results show that incorporation of CDEA in place of betaine gives little or no benefit as regards lather, and can be detrimental.

Example 7

25 The procedure of Example 3 was repeated using solutions of cocoyl isethionate (Fenopon AC78 as in Examples 1 and 3) coconut amidopropyl betaine (Rewoteric AMB14 as in Example 2) and a third detergent in a weight ratio of 50:40:10 by weight. The third detergent was

selected from:

- (i) sodium lauryl sarcosinate (Hamposyl L-95 from W.R. Grace)
- (ii) sulphosuccinate (Rewopol SBFA as in Example 1)
- 5 (iii) sodium lauryl sulphate (Empicol LX 28 from Albright and Wilson)
- (iv) alcohol ethoxylate (C_{12-13} fatty alcohol ethoxylated with average 8 ethylene oxide residues).

10 All solutions contained 15% by weight detergent. A control solution as in Examples 3 and 4 was used. The results were:

Table 7

		<u>Total panel scores</u>	
		<u>lather volume</u>	<u>lather creaminess</u>
15	Third detergent		
	(i) sarcosinate	110	101
	(ii) sulphosuccinate	102	97
	(iii) lauryl sulphate	91	106
	(iv) alcohol ethoxylate	82	90
20	Control 86 SLES:14 betaine	94	103

This shows that the alcohol ethoxylate had an adverse effect on lather which the anionic detergents did not.

25 Example 8

A number of aqueous solutions of sodium cocoyl isethionate and a second detergent were made and assessed for lathering as in Example 3. The isethionate was

Fenopon AC78 as in Examples 1 and 3, coconut amidopropyl sulphobetaine (Rewoteric AM-CAS as in Example 1) or coconut amidopropyl betaine (CAPB) derived from middle cut coconut oil so that substantially all the long chain acyl groups

5 R¹CO contain 12 or 14 carbon atoms (Tegobetain L5351 from Goldschmidt). The same ether sulphate/betaine control solution was used. All solutions contained 15% by weight of detergent. The results obtained were:

10

Table 8

<u>proportions</u>			<u>panel scores</u>	
isethionate	sulphobetaine	CAPB	lather volume	creami-ness
70	30	--	106	94
15 60	40	--	113	106
50	50	--	97	88
70	--	30	95	100
60	--	40	96	111
50	--	50	111	108
20 Control	86 SLES: 14 betaine		95	93

This demonstrates the effectiveness of these two zwitterionic detergents.

25

Example 9

The procedure of Example 3 was repeated using two solutions of cocoyl isethionate (Fenopon AC78 as in Examples 1 and 3), coconut amidopropyl betaine (Rewoteric AMB14 as in Example 2) and in one solution sodium lauryl

ether sulphate with average 3EO (Empicol 0251 as in Example 1). The solutions were thickened by incorporation of ammonium chloride.

- 5 One solution contained isethionate, CAPB and SLES in a weight ratio of 33:54:13. The other contained equal weights of isethionate and CAPB without SLES. Both contained 15% by weight detergent in total. Both were isotropic.

10

The panel scores for lather volume and creaminess were the same for both solutions and matched the scores for 86 SLES: 14 betaine used as control.

15 Example 10

Various compositions were assessed for mildness by the zein test as in Example 2. The compositions contained sodium cocoyl isethionate (Fenopon AC78 as in Example 1) sodium lauryl ether sulphate with average 3EO and CAPB, CDEA or a mixture of the two as in Example 5.

- 20 All solutions contained 3% detergent. The proportions of the constituents, as percentages by weight of total detergent, and the zein solubilisation figures were as follows:
- 25

Table 9

% by weight of total detergent

Isethionate	50	50	50	50	50	50	33	33	33	33	16	16	16
CAPB	40	-	20	25	-	54	-	27	33	-	42	-	21
CDEA	-	40	20	-	25	-	54	27	-	33	-	42	21
SLES	10	10	10	25	25	13	13	13	33	33	42	42	42

Zein solubilisation

(%N)

0.15 0.38 0.31 0.25 0.49 0.07 0.27 0.20 0.19 0.40 0.07 0.34 0.29

These results show very clearly that alkanolamide is detrimental to mildness even when used jointly with CAPB.

5 Example 11

Binary mixtures were tested for mildness by zein test, using solutions containing 10% by weight detergent. The mixtures contained sodium cocoyl isethionate with CAPB, a simple betaine (Empigen BB) or CDEA. The results obtained

10 were:

Table 10

		<u>% by weight of total detergent</u>								
15	Isethionate	70	70	70	60	60	60	50	50	50
	CAPB	30	-	-	40	-	-	50	-	-
	Betaine	-	30	-	-	40	-	-	50	-
	CDEA	-	-	30	-	-	40	-	-	50
Zein solubalisation (%N)		0.52	0.49	1.01	0.34	0.34	0.85	0.24	0.21	0.66

20

This data shows that there is very little difference in mildness between CAPB and simple betaine. It confirms that ethanolamide is detrimental to mildness.

Example 12

A shower gel contained the following:

		<u>% by weight</u>
5	Sodium cocoyl isethionate (Fenopon AC78)	5.0
	Coconut amidopropyl betaine (Tegobetain L7)	8.0
10	Sodium lauryl ether sulphate 3EO (Empicol 0251)	2.0
	Isopropyl palmitate	0.5
15	Opacifier	1.0
	Glycerol	1.25
20	Preservative	0.07
	Sodium chloride	3.5
	Perfume	1.5
25	Water	balance to 100%

The three detergents were fully dissolved in the aqueous phase. The opacifier, a polystyrene latex, was in suspension.

Example 13

A hand washing composition contained the following:

	<u>% by weight</u>
5 Sodium cocoyl isethionate	2.0
Coconut amidopropyl betaine	5.3
Sodium lauryl ether sulphate 3EO	5.3
Ethylene glycol monostearate	2.0
Sodium chloride	2.0
10 Triclosan (antimicrobial agent)	0.4
Perfume	0.4
Preservatives, colourants	q.s.
Water	balance to 100%

- 15 The three detergents were fully dissolved in the aqueous phase. The ethylene glycol monostearate was in suspension and functioned as an opacifier.

Example 14

A shower gel contained the following:

		<u>% by weight</u>
5	Sodium cocoyl isethionate (Fenopon AC78)	7.5
	Coconut amidopropyl betaine (Rewoteric AMB14)	5.0
10	Sodium lauryl ether sulphate 3EO (Genapol ZRO)	2.5
	Preservative	0.07
15	Ammonium chloride	1.8
	Perfume	1.0
20	Water	balance to 100%

This is a clear gel.

Example 15

A hair shampoo contained the following:


		<u>% by weight</u>
5	Sodium cocoyl isethionate (Fenopon AC78)	5.0
	Coconut amidopropyl betaine (Rewoteric AMB14)	5.0
10	Sodium lauryl ether sulphate 3EO (Genapol ZRO)	5.0
	Preservative	0.07
15	Sodium chloride	2.2
	Perfume	0.5
20	Water	balance to 100%

The shampoo was a clear, isotropic liquid.

CLAIMS:

1. A detergent composition in the form of an aqueous
5 liquid or gel comprising 10% to 50% by weight of a
detergent mixture which comprises

(a) 10 to 60% by weight of the detergent mixture of
a fatty acyl isethionate of formula



10 where R is an alkyl or alkenyl group of 7 to 21
carbon atoms and M is a solubilising cation such as
sodium, potassium, ammonium or substituted ammonium;

(b) 10 to 80% by weight of the detergent mixture of
a zwitterionic detergent which has a hydrophilic head
15 group containing a quaternary nitrogen atom and at least
one acid group,

(c) 10 to 55% by weight of the detergent mixture of
a further anionic detergent,

wherein the amount by weight of the fatty acyl
20 isethionate (a) is not more than three times the amount by
weight of the zwitterionic detergent (b), the total of (a)
and (b) is from 45 to 90% by weight of the detergent
composition, and the composition is sufficiently free of
alkanolamide detergents that the amount by weight of
25 alkanolamide is not more than one quarter the amount of
the zwitterionic detergent (b).

2. A composition according to claim 1 wherein the

zwitterionic detergent (b) is 30% to 60% by weight of the detergent mixture.


3. A composition according to claim 1 or claim 2 wherein
5 the fatty acyl isethionate (a) and zwitterionic detergent (b) together constitute 55 to 75% by weight of the detergent mixture.

4. A composition according to claim 1, claim 2 or claim
10 3 wherein the weight ratio of the fatty acyl isethionate and zwitterionic detergent lies in a range from 2:1 to 1:2.

5. A composition according to any one of claims 1 to 4
15 containing at least 6% by weight of the whole composition of the fatty acyl isethionate (a) and at least 3% by weight of the whole composition of the zwitterionic detergent (b).

20 6. A composition according to any one of claims 1 to 5 wherein at least three quarters of the groups R of the acyl isethionate are alkyl of 11 to 17 carbon atoms.

7. A composition according to any one of claims 1 to 6
25 wherein the zwitterionic detergent is of the formula



where R^1 is alkyl or alkenyl of 7 to 18 carbon atoms

R^2 and R^3 are each independently alkyl, hydroxyalkyl
or carboxyalkyl of 1 to 3 carbon atoms

m is 2 to 4

5 n is 0 or 1.

8. A composition according to claim 7 wherein at least
three quarters of the groups R^1 have 12 to 14 carbon atoms
and R^2 and R^3 are methyl.

10

9. A composition according to any one of claims 1 to 8
wherein the further anionic detergent (c) is alkyl ether
sulphate with an average of at least 2.5 ethylene oxide
residues per molecule.

15

10. A composition according to any one of claims 1 to 9
which also contains inorganic electrolyte.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 91/01955

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 A61K7/08; C11D1/12; C11D1/94		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	A61K ; C11D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	WO,A,9 012 860 (UNILEVER PLC / UNILEVER NV) 1 November 1990 see the whole document	1-10
A	GB,A,2 114 996 (COLGATE-PALMOLIVE COMPANY) 1 September 1983 see the whole document	1-10
A	BE,A,840 667 (JOHNSON & JOHNSON) 12 October 1976 see page 6, line 1 - page 7, line 4; claims 1-12	1-10
A	EP,A,0 207 642 (DEUTSCHE ICI GMBH) 7 January 1987 see column 3, line 54 - line 58; claims 1-9	1-10
A	EP,A,0 277 876 (LABORATOIRE LACHARTRE SOCIETE ANONYME) 10 August 1988 see page 3, line 24 - page 4, line 60; claims 1-12	1-10
-/-		
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
20 FEBRUARY 1992	12.03.92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	Els Vonk	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	US,A,3 723 356 (ARNO CAHN ET AL.) 27 March 1973 see the whole document ---	1-10

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. GB 9101955
SA 53066**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 20/02/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9012860	01-11-90	US-A- 4954282	04-09-90
		AU-A- 5527490	16-11-90
		CA-A- 2014831	19-10-90
		EP-A- 0469029	05-02-92
GB-A-2114996	01-09-83	US-A- 4554098	19-11-85
		AU-B- 559168	26-02-87
		AU-A- 1164383	25-08-83
		CA-A- 1205347	03-06-86
		CH-A- 654590	28-02-86
		DE-A- 3305197	01-09-83
		FR-A,B 2522012	26-08-83
		SE-B- 456425	03-10-88
BE-A-840667	12-10-76	None	
		None	
EP-A-0207642	07-01-87	None	
EP-A-0277876	10-08-88	FR-A- 2610194	05-08-88
US-A-3723356	27-03-73	DE-A- 1617229	18-02-71
		FR-A- 1535087	
		GB-A- 1193469	03-06-70
		NL-A- 6711312	19-02-68
		SE-A,B 356530	
		US-A- 3649543	14-03-72